

Publikujeme v zahraničí

Onkológia (Bratisl.), 2020;15(2):148

SUPORTÍVNA LIEČBA

Gudiol C, Albasanz-Puig A, Laporte-Amargós J, Pallarès N, Mussetti A, Ruiz-Camps I, Puerta-Alcalde P, Abdala E, Oltolini C, Akova M, Montejo M, Mikulska M, Martín-Dávila P, Herrera F, Gasch O, **Drgona L**, Paz Morales H, Brunel AS, García E, Isler B, Kern WV, Morales I, Maestro-de la Calle G, Montero M, Kanj SS, Sipahi OR, Calik S, Márquez-Gómez I, Marin JI, Gomes MZR, Hemmatti P, Araos R, Peghin M, Del Pozo JL, Yáñez L, Tilley R, Manzur A, Novo A, Carratalà J; IRONIC study group.

A clinical predictive model of multidrug resistance in neutropenic cancer patients with bloodstream infection due to *Pseudomonas aeruginosa* (IRONIC study).

Antimicrob Agents Chemother. 2020 Feb 3. pii:AAC.02494-19.

Background: We aimed to assess the rate and predictive factors of bloodstream infection (BSI) due to multidrug-resistant (MDR) *Pseudomonas aeruginosa* (PA) in neutropenic cancer patients.

Methods: We performed a multicenter, retrospective cohort study including onco-hematological neutropenic patients with BSI due to PA conducted across 34 centers in 12 countries from January 2006 to May 2018. A mixed logistic regression model was used to estimate a model to predict multidrug resistance of the causative pathogens.

Results: Of a total of 1217 episodes of BSI due to PA, 309 episodes (25.4%) were caused by MDR strains. The rate of multidrug resistance increased significantly over the study period ($p=0.033$). Predictors of MDRPA BSI were prior therapy with piperacillin/tazobactam (odds ratio [OR], 3.48; 95% confidence interval [CI], 2.29–5.30), prior antipseudomonal carbapenem use (OR, 2.53; 95% CI, 1.65–3.87), fluoroqu-

inolone prophylaxis (OR, 2.99; 95% CI, 1.92–4.64), underlying hematological disease (OR, 2.09 95% CI, 1.26–3.44) and the presence of a urinary catheter (OR, 2.54; 95% CI, 1.65–3.91), whereas older age (OR, 0.98; 95% CI, 0.97–0.99) was found to be protective.

Conclusions: Our prediction model achieves good discrimination and calibration, thereby identifying neutropenic patients at higher risk of BSI due to MDRPA. The application of this model using a web-based calculator may be a simple strategy to identify high-risk patients, who may benefit from the early administration of a broad-spectrum antibiotic coverage against MDR strains according to the local susceptibility patterns, thus avoiding the use of broad-spectrum antibiotics in patients at low risk of resistance.

ABSTRAKTY A PRÍSPEVKY Z KONFERENCIÍ

GENITOURINÁRNE MALIGNITY

Hapakova N, Chovanec M, Rejlekova K, Kalavska K, Obertova J, Palacka P, De Angelis V, Sycova-Mila Z, Mardiak J, Mego M.

Effect of prophylactic anticoagulation on incidence of venous thromboembolism in testicular germ cell tumor patients

J Clin Oncol 38, 2020 (suppl 6; abstr 422)

Rejlekova K, Kalavska K, Celec P, Makovnik M, Chovanec M, Hapakova N, De Angelis V, Obertova J, Palacka P, Sycova-Mila Z, Mardiak J, Mego M.

Biomarkers of lung damage with possible predictiveness of ARDS within CS in super high-risk patients with germ cell tumors

J Clin Oncol 38, 2020 (suppl 6; abstr 411)

KARCINÓM PLÚC

Kasan P, Berzinec P, Cerna M, Andrasina I, Cipkova A, Chowaniecova G, Denkova L, Godal R, Kuzmova H, Svihelova-Liskova Z, Martak M, Mazal J, Obertova J, Pribulova Z, Urda M, Vesela M.

First experience with atezolizumab in Slovakia

IASLC 2019 World Conference on Lung Cancer September 7–10, 2019, Barcelona, Spain

Berzinec P, Kasan P, Mazal J, Andrasina I, Cipkova A, Chowaniecova G, Martak M, Urda M, Vesela M, Jancokova I, Kuliskova I, Plank L.

Advanced nsclc treated with gefitinib or erlotinib for five years or longer – updated data from the retrospective slovakian study

IASLC 2019 World Conference on Lung Cancer September 7 – 10, 2019, Barcelona, Spain

Berzinec P, Kasan P, Mazal J, Andrasina I, Cipkova A, Bodorova M, Demo P, Chowaniecova G, Jancokova I, Kuliskova I, Martak M, Palkovacsova I, Urda U, Vesela M, Plank L.

Five-year progression free survival in patients with advanced NSCLC treated with EGFR-TKIs

18th Central European Lung Cancer Conference, Budapest, 21 – 23 November 2019

KARCINÓM PRSNÍKA

Mego M, Kalavska K, Karaba M, Minarik G, Benca J, Sedlackova T, Manasova D, Pindak D, Mardiak J, Cierna Z.

CTC with EMT phenotype are associated with PD-L1 expression in tumor associated stroma in primary breast cancer patients.

Cancer Research, Vol. 80, Supplement 1 (2020), s. 1-1. San Antonio Breast Cancer Symposium, San Antonio, TX, USA, December 10 – 14, 2019